

FINAL PROGRESS REPORT FOR NAS5-98191:
A THREE-DIMENSIONAL ANALYSIS OF THE GALACTIC GAMMA-RAY EMISSION
RESULTING FROM COSMIC-RAY INTERACTIONS WITH THE
INTERSTELLAR GAS AND RADIATION FIELDS

TASK LEADER: Thomas J. Sodroski
ATR: Eli Dwek

Description of Activities:

The contractor will provide support for the analysis of data under ADP (NRA 96-ADP-09; Proposal No . 167-96adp). The primary task objective is to construct a 3-D model for the distribution of high-energy (20 MeV - 30 GeV) gamma-ray emission in the Galactic disk. Under this task the contractor will utilize data from the EGRET instrument on the Compton Gamma-Ray Observatory, H I and CO surveys, radio-continuum surveys at 408 MHz, 1420 MHz, 5 GHz, and 19 GHz, the COBE Diffuse Infrared Background Experiment (DIRBE) all-sky maps from 1 to 240 μm , and ground-based B, V, J, H, and K photometry. The respective contributions to the gamma-ray emission from cosmic ray/matter interactions, inverse Compton scattering, and extragalactic emission will be determined.

This task is supported by T. J. Sodroski (EITI).

Significant Accomplishments:

During this second year of our 2-year program we have successfully derived the respective contributions to the high-latitude ($|b| > 20^\circ$) gamma-ray emission from cosmic-ray/matter interactions, inverse Compton scattering (i.e., including the Galactic gamma-ray halo component), and the extragalactic gamma-ray background component. A collaboration with the EGRET team at Goddard was established for continuation of project.

We were provided with source-subtracted EGRET data at high Galactic latitudes, derived using the source-removal algorithm developed by the EGRET team. A map of H-atom column density covering the total sky was obtained from the Leiden/Dwingeloo survey of H I over the northern sky ($\delta > -30^\circ$) by Hartmann & Burton (1993, 1997), and the Cleary, Heiles, and Haslam (1979) H I survey of the southern sky ($\delta < -30^\circ$). Maps of the inverse Compton emission in each EGRET energy interval were generated using the 3-D map of the Galactic interstellar radiation field (ISRF) that was derived by our group from our analysis of COBE/DIRBE data and ground-based B, V, J, H, and K photometry of the Milky Way (See Final Progress Report For S-92504-Z). The local cosmic-ray electron intensity distribution of Fichtel et al. (1991), modified by Bertsch et al. (1993), was adopted. It was further assumed that the radial gradient of the CR electron intensity is equal to the gamma-ray emissivity gradient derived by Strong et al. (1988; Case 3, where shape of radial distribution is independent of gamma-ray energy), which is consistent with the results of Strong & Mattox (1996) using EGRET data. We set up an idl algorithm for generating maps of the source-subtracted gamma-ray emission, H I emission, and Inverse Compton model emission at comparable resolution.

We implemented a modelling algorithm to perform the high latitude decomposition with Poisson statistics. The algorithm was fitted to the point source-subtracted EGRET data for all 10 bands simultaneously (10480 lines of sight) in order to obtain the value of the absolute scaling factor for the local cosmic-ray electron intensity distribution, and for each EGRET energy interval the value of the gamma-ray emissivity due to nuclear interactions and bremsstrahlung collisions, the scale height of the cosmic-ray electrons responsible for the inverse Compton emission, and the extragalactic background intensity (total of 31 parameters). Our results are presented in Figures 1 – 3.

The determination of the contribution of inverse Compton scattering to the Galactic gamma-ray emission at low latitudes is currently being done in collaboration with the EGRET team at Goddard. We have reformatted our Galactic interstellar radiation field (ISRF) determination onto an interpolated grid suitable for their Galactic plane models (Bertsch et al. 1993; Hunter et al. 1997), and have been in close communication with their group to verify that our model is incorporated correctly.

Three papers (Sodroski et al. 2000; Odegard et al. 2000; Hunter et al. 2000) discussing the details of the analysis and results of our two year program are currently in progress. Sodroski et al. (2000) describe the analysis and results of the decomposition of the high-latitude EGRET data. Odegard et al. (2000) discuss the derivation of the local ISRF using DIRBE data. The results of the decomposition of the low-latitude EGRET data will be presented by Hunter et al. (2000).

References:

- Bertsch, D. L., et al. 1993, ApJ, 416, 587
 Cleary, M. N., Heiles, C., & Haslam, C. G. T. 1979, A&AS, 36, 95
 Fichtel, C. E., Özel, M. E., Stone, R. G., & Sreekumar, P. 1991, ApJ, 374, 134
 Hartmann, D., & Burton, W. B. 1993, in AIP Conf. Proc. 278, Back to the Galaxy, ed. F. Verter
 & S. S. Holt (New York: AIP), 279
 Hartmann, D., & Burton, W. B. 1997, Atlas of Galactic Neutral Hydrogen (Cambridge:
 Cambridge University Press)
 Hunter, S. D., et al. 1997, ApJ, 481, 205
 Hunter, S. D., et al. 2000, in progress
 Odegard, N. P., et al. 2000, in progress
 Sodroski, T. J., et al. 2000, in progress
 Strong, A. W., et al. 1988, A&A, 207, 1
 Strong, A. W., & Mattox, J. R. 1996, A&A, 308, L21




Fig. 1. — Emissivity spectrum due to nuclear interactions and bremsstrahlung collisions.




Fig. 2. — The distribution of the scale height of the cosmic-ray electrons responsible for the inverse Compton emission in each EGRET energy interval.




Fig. 3. — The extragalactic diffuse emission spectrum.

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 5, 2000	3. REPORT TYPE AND DATES COVERED Final: September 1998 – June 2000	
4. TITLE AND SUBTITLE A Three-Dimensional Analysis of the Galactic Gamma-Ray Emission Resulting from Cosmic-Ray Interactions with the Interstellar Gas and Radiation Fields			5. FUNDING NUMBERS NAS5-98181	
6. AUTHORS Dr. Thomas Sodroski				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Space Applications Corporation 9315 Largo Drive West, Suite 250 Largo, MD 20774			8. PERFORMING ORGANIZATION REPORT NUMBER 1042-006/FINAL	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA/Goddard Space Flight Center Code 216 Greenbelt, MD 20771			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The contractor will provide support for the analysis of data under ADP (NRA 96-ADP-09; Proposal No. 167-96adp). The primary task objective is to construct a 3-D model for the distribution of high-energy (20 MeV - 30 GeV) gamma-ray emission in the Galactic disk. Under this task the contractor will utilize data from the EGRET instrument on the Compton Gamma-Ray Observatory, H I and CO surveys, radio-continuum surveys at 408 MHz, 1420 MHz, 5 GHz, and 19 GHz, the COBE Diffuse Infrared Background Experiment (DIRBE) all-sky maps from 1 to 240 μ m, and ground-based B, V, J, H, and K photometry. The respective contributions to the gamma-ray emission from cosmic ray/matter interactions, inverse Compton scattering, and extragalactic emission will be determined.				
14. SUBJECT TERMS Galaxy - diffuse radiation - cosmic rays - gamma rays			15. NUMBER OF PAGES 4	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	